

Non-odontogenic Cysts

Dr. Ioannis G. Koutlas
Division of Oral Pathology

All pictures are intellectual property of the Division of Oral and Maxillofacial Pathology or its Faculty. Duplication or any unauthorized use is prohibited.

Developmental Cysts

- a.k.a. fissural cysts
- Exact pathogenesis of some of them uncertain
- Generally, slow increase
- May be identified incidentally

Palatal and gingival cysts of newborns

- Common (more than half of neonates)
- Small, occasionally in clusters
- Embryogenesis
 - Entrapped epithelium during formation of 2^o palate
 - Epithelial remnants of salivary glands
- Epstein pearls: midline
- Bohn's nodules: hard/soft palate
- Gingival cysts: keratin filled
- No treatment, spontaneous rupture




Nasopalatine duct cyst

- a.k.a. incisive canal cyst, nasopalatal cyst
- Most common non-odontogenic cyst
- Embryogenesis
 - Remnants of nasopalatine duct
 - Incisive canals exit via a common foramen
 - Rarely two foramina = canals of Scarpa



Nasopalatine duct cyst

- 4th to 6th decades
- Swelling, drainage, pain or asymptomatic
- Radiolucency (sometimes heart-shaped) at or near the midline, between and apical to central incisors
- Incisive canal size ~ 6mm
- Cysts may reach 3 cm and cause a “through-and-through” expansion of the jaw



Nasopalatine duct cyst

- Histology
 - Stratified squamous epithelium
 - Pseudostratified columnar epithelium
 - Simple epithelium
 - Respiratory epithelium
 - Nerve bundles and muscular arteries and veins
 - Cartilagenous remnants
- Treatment
 - Surgical enucleation

Median palatal (palatine) cyst

- Most cases are posteriorly positioned nasopalatine cysts
- Fusion between palatal shelves of the maxilla
- Midline of hard palate
- Swelling (it's a must) ~ 2cm
- Well circumscribed radiolucency

Nasolabial cyst

- Upper lip lateral to midline
 - Medial - lateral nasal and maxillary processes OR
 - Misplaced epithelium of nasolacrimal gland
- Elevation of nasal ala
- No pain except if infected
- More women than men
- Can be bilateral
- Surgical removal

“Median mandibular cyst”

- Controversial
 - There are no epithelial lined processes
- Midline of mandible
- Most of them periapical or lateral periodontal cysts or odontogenic keratocysts




“Globulomaxillary cyst”

- Controversial
 - Alleged fusion of maxillary process and medial nasal process
 - No epithelial entrapment
 - Most if not all cysts are of odontogenic origin
- Lateral incisor and canine
- Surgical excision



Epidermoid cyst

- a.k.a. infundibular cyst, sebaceous cyst (wrong term); epidermal inclusion cyst (after trauma)
- More frequent in acne-prone areas
- DERIVE FROM HAIR FOLLICLE
- Unusual before puberty except when associated with Gardner syndrome
- Nodular, fluctuant subcutaneous lesion, white or yellow
- Cavity lined by epithelium containing keratin



Dermoid cyst

- Benign cystic teratoma (tissue form all three embryonic layers)
- Dermoid cyst is a forme fruste
- Intraoral epidermoid cyst
- Midline of floor of mouth or rarely displaced laterally – above the geniohyoid muscle
- Submental swelling – below the geniohyoid muscle
- Few mm to several cm
- Children and young adults
- Doughy or rubbery mass; may drain



Thyroglossal duct (tract) cyst

- Review thyroid development
- Remnants of the thyroglossal duct epithelium
- Have been described in families
- Midline (foramen cecum → suprasternal notch)
 - Except if they are located in the area of thyroid cartilage
- More frequently below the hyoid
- ! Base of tongue → airway obstruction
- Clue: If it maintains attachment to hyoid bone or tongue it will move vertically during swallowing or protrusion of the tongue; can cause fistulous tract
- Thyroid tissue may be absent histologically



Branchial cleft cyst


- a.k.a. cervical lymphoepithelial cyst
- From branchial clefts (2nd arch), or cystic change of parotid gland epithelium entrapped in upper cervical lymph nodes (not valid)
- Lateral upper neck along the anterior border of the sternocleidomastoid muscle
- Young adults, fluctuant, between 1-10 cm, more frequently on the left
- Occasionally a fistula
- Contain lymphoid tissue (occasionally not though)
- Rare examples of malignant transformation



Oral lymphoepithelial cyst

- Invaginations of epithelium resulting in pouches or crypts that may fill with keratin debris
- Salivary gland or surface epithelium that becomes entrapped
- Less than 1 cm
- Firm or soft to palpation
- White or yellow
- DD: lipoma






Antral pseudocysts

- Antral pseudocyst
- Sinus mucocele
 - Surgical ciliated cyst
 - Obstruction (ostium) mucocele
- Retention cyst
 - Obstruction of ducts of seromucous glands
 - Invagination of epithelium

Antral pseudocyst

- Common finding of mostly panoramic radiographs
- Dome-shaped, slightly radiopaque
- Floor of maxillary sinus
- NOT A CYST
- Inflammatory exudate; probable odontogenic origin
- Other causes may be: allergy, sinus infection
- No treatment necessary if not associated with signs and symptoms (enlargement; headache)
- EVALUATION OF TEETH





Sinus mucocele

- Whole sinus is cloudy when ostium obstructed
- Post surgical cysts can enlarge
- Surgical removal or meatal antrostomy

Retention cyst

- Very small to be detectable